

Drug Facts		Drug Facts (continued)	
Active ingredient (in each tablet)		Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Aspirin 81 mg (NSAID)*	Purpose Pain reliever	Directions ■ drink a full glass of water with each dose	
*nonsteroidal anti-inflammatory drug		■ adults and children 12 years of age and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor	
Uses		■ children under 12 years of age: do not use unless directed by a doctor	
<ul style="list-style-type: none"> ■ temporarily relieves minor aches and pain ■ ask your doctor about other uses for St. Joseph Safety Coated 81 mg Aspirin 		Other information	
Warnings		<ul style="list-style-type: none"> ■ store at 20-25°C (68-77°F) ■ close cap tightly after use 	
<p>Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Aspirin may cause a severe allergic reaction, which may include:</p> <ul style="list-style-type: none"> ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you</p> <ul style="list-style-type: none"> ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed 		Inactive ingredients FD&C red #40, FD&C yellow #6, methacrylic acid copolymer, microcrystalline cellulose, pregelatinized starch, silicon dioxide, sodium bicarbonate, sodium lauryl sulfate, stearic acid, talc, triethyl citrate	
<p>Do not use if you are allergic to aspirin or any other pain reliever/fever reducer</p> <p>Ask a doctor before use if</p> <ul style="list-style-type: none"> ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma <p>Ask a doctor or pharmacist before use if you are taking a prescription drug for gout, diabetes or arthritis</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ ringing in the ears or loss of hearing occurs ■ redness or swelling is present ■ new symptoms occur. These could be signs of a serious condition. <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p>		Questions or comments? Call 1-855-785-6381 weekdays 9 AM to 5 PM EST	